

Degrees of residence of Guiana dolphins (*Sotalia guianensis*) in Ilha Grande Bay, south-eastern Brazil: a preliminary assessment

MARIANA DE A. ESPÉCIE^{1,2}, RODRIGO H.O. TARDIN^{1,2} AND SHEILA M. SIMÃO²

¹Programa de Pós-Graduação em Biologia Animal, Universidade Federal Rural do Rio de Janeiro, Rodovia BR 465, km 7, Seropédica, Rio de Janeiro, 23890-000, Brazil, ²Laboratório de Bioacústica e Ecologia de Cetáceos, Departamento de Ciências Ambientais, Universidade Federal Rural do Rio de Janeiro, Rodovia BR 465, km 7 Seropédica, Rio de Janeiro, 23890-000, Brazil

The Guiana dolphin, Sotalia guianensis, is a small delphinid found in coastal areas from northern Honduras to southern Brazil. Little is known about the ecology of this species in several areas of its geographical distribution. In this paper, we present new data about the residency of Guiana dolphins in Ilha Grande Bay, south-eastern Brazil. Boat surveys were conducted at the study area from May 2007 to March 2008. Applying the photo-identification technique, we took pictures of dolphins' dorsal fins, looking for natural markings on them. A total of 17,969 photographs were taken, from which was created a databank of 462 distinct dolphins. Individuals showed different degrees of residency in the bay. More than 50% of the catalogued dolphins were considered non-residents and/or showed a low degree of residence. The average interval between resightings was 35 ± 27.4 days. Animals without markings (calves and juveniles) accounted for 45% ($N = 2917$) of photographs taken. These results indicate that this population has a fluid structure including different individuals over time. We suggest a possible relationship between the presence of some individuals and local resource availability, such as prey and protected areas.

Keywords: Guiana dolphin, photo-identification, degrees of residence, Ilha Grande Bay

Submitted 28 February 2010; accepted 4 June 2010; first published online 21 September 2010

INTRODUCTION

As a primary tool for assessing some population parameters, the use of the photo-identification technique has improved our knowledge about the Guiana dolphin, *Sotalia guianensis* (van Bénédén, 1864), in its natural habitat (Flores, 1999; Pizzorno, 1999; Simão *et al.*, 2000; Santos *et al.*, 2001; Rossi-Santos *et al.*, 2007). Using a dataset composed of photographs of dolphins' dorsal fins, it is possible to distinguish each individual based on natural markings (Würsig & Würsig, 1977; Hammond *et al.*, 1990). One of the ecological parameters that can be assessed through a photo-ID catalogue is residency, which refers to the amount of time that a certain individual spends in a particular geographical area (Wells & Scott, 1990; Wells, 1991).

Of the entire geographical distribution of this species, the largest aggregation of Guiana dolphins is reported to be located in western Ilha Grande Bay ($23^{\circ}02'S$ $44^{\circ}26'W$) (Lodi & Hetzel, 1998). However, data about many population features are scarce, which makes any conservation decision affecting the species in the area difficult. Ilha Grande Bay is an area of high primary productivity, which guarantees the sustainability of several marine species that inhabit it (Nogara, 2000). Previous studies in the area have shown

that Guiana dolphins use the bay mainly for feeding purposes and as a care giving area (Lodi, 2003a, b).

Food resources and protection from potential predators are considered to be the main factors that influence the distribution and permanence of a species in a certain area (Mazzoil *et al.*, 2008). Estuaries, such as Ilha Grande Bay, are usually considered to contain large concentrations of nutrients that support a great number of filter-feeding zooplankton and fish (Ballance, 1992). Guiana dolphins may inhabit this bay mainly because of the potential prey abundance that this area supports. Early studies indicated that the area is predator-free, representing a valuable place for caring for and teaching young offspring (Lodi, 2003a, b). However, with the increase of anthropogenic activities in the area, such as overfishing and boat traffic, as well as high levels of pollutants and contaminants, the natural stability of all species that inhabit this ecosystem, including *S. guianensis*, may become disturbed. The main objective of this work was to characterize the degree of residence of Guiana dolphins in Ilha Grande Bay, Brazil. The results provided by this study represent the first step in acquiring knowledge of diverse characteristics of *S. guianensis* in the area.

MATERIALS AND METHODS

Study area

Ilha Grande Bay is located on the southern coast of Rio de Janeiro State, Brazil ($23^{\circ}02'S$ $44^{\circ}26'W$). According to its

Corresponding author:

M.A. Espécie

Email: mariana_especie@ufrjr.br

geographical, hydrological and physiographic aspects, it is divided into three different units: the east side, the central part and the west side (Mahiques, 1987). The present study took place in a delimited area of 190 km² on the west side of Ilha Grande Bay (Figure 1), which is characterized by shallow depths (~10 m on average). Previous studies have shown that Guiana dolphins are found mainly in this part of the bay (Lodi & Hetzel, 1998; Lodi, 2003a, b). The marine environment of Ilha Grande Bay's west side is considered a transition area between mainland and sea (Nogara, 2000). An organic matter influx from the continent takes place due to water input from continental rivers and from mangrove swamps around the bay. According to Signorini (1980), during the summer, the west side of the bay receives nutrients from deep water (southern Atlantic central waters) that enters the bay by its west channel, causing an upwelling phenomenon. The area presents a great diversity of microhabitats, such as islands, mangrove swamps, rocky cliffs, muddy areas and sandy beaches (Lodi, 2003a). At the same time, the study area is of great economic interest, with different anthropogenic activities being developed around it.

Data collection

This research was conducted from May 2007 to March 2008 during all four seasons of the year, using a 7.5-m vessel (22 HP) that cruised at 4 knots along the west side of Ilha Grande Bay. Boat trips were performed only when climatic conditions were favourable for dolphin sightings (i.e. without rain and sea state <3). The boat travelled without any systematic route inside the delimited area until one or more groups of dolphins were sighted by experienced cetacean observers, who recorded the animals' geographical position using a GPS, time of sighting, group size and behaviour. Groups were approached to take the photographs with the boat at slow speed and keeping a physical distance of about 10 m along a parallel route. All photographs were taken using a digital camera (Canon EOS 20D Digital fitted with Canon Zoom EF 75-300 mm lens) and a 1 GB memory card. The photographs were taken by the same observer during the entire study period. The sampling protocol was to photograph as many animals as possible in each school. Field effort ended when there was no more space on the memory card and/or sudden climate changes took place.

Data analysis

Data gathered by the GPS device and the digital camera were transferred to computers for further laboratory analysis. All photographs were passed through an extensive sorting and selection process to prevent errors during the cataloguing process (Gunnlaugsson & Sigurjónsson, 1990; Stevick *et al.*, 2001). For this reason, two experienced observers selected only pictures of 'good' or 'excellent' quality (well focused, with enough light and little blurring) during this stage. Nicks and notches along the dorsal fins were the main features used to distinguish individual dolphins (Würsig & Würsig, 1977), but in many cases, scars and injuries on dolphins' bodies were used as auxiliary marks for identification.

We adopted the methodology proposed by Mazzoil *et al.* (2004) for sorting and selecting the digital photographs. Matches with previously identified individuals were made by visually comparing each new photograph with all others in

an existing library of dorsal fin pictures. Animals that could not be matched received a new identification number. The best picture of each was added to the digital photo-ID catalogue. Animals without markings on their dorsal fins were classified as juveniles or calves, based on their appearance. Dolphins that had one-half to three-quarters of an adult body length were defined as juveniles. Small dolphins with no more than one-quarter of an adult body length and with pinkish-grey coloration were assumed to be calves. As young dolphins have no distinctive marks, they were not included in the catalogue.

A Pearson correlation (r_p) was used to test whether our sample effort was representative. For this reason, sample standardization was performed by calculating the proportions of captured and recaptured individuals in each survey compared to the cumulative number of captured individuals (capture and recapture rates, respectively). Definitions used to determine residency in this paper are those stated by Ballance (1990). Individuals seen only one time were considered non-residents. Resident dolphins were sighted more than once, and the term is used in a relative sense; that is, one animal can be said to exhibit a higher or lower degree of residence than another. Among resident dolphins, we observed three different aspects of residence: the number of times an animal was seen in the area; the time interval between the first and last sightings of each individual; and the average number of days between sightings of each dolphin. Analysing these three items together, we could determine the degree of residence (e.g. high, medium or low) of each individual. Photographs depicting individuals without distinctive markings on their dorsal fins (calves and juveniles) were used to estimate the proportion of immature individuals in the population.

RESULTS

Twenty-three boat surveys were conducted in Ilha Grande Bay during all seasons of the year. Dolphins were seen in 17 surveys (73.9%) and were observed to be widely distributed in the study area (Figure 1). A total of 17,969 photographs were taken, of which 6014 pictures (33.5%) were characterized as being of 'good' or 'excellent' quality. Of these, about 45% (N = 2917) pictured animals without markings (calves and juveniles). The mean number of pictures taken in each survey was 1057 (± 598). A total of 1043 subgroups were sighted, and the average subgroup size was 18.7 dolphins (± 19.7). Group sizes in each survey ranged from 20 to 350 dolphins (mean 165 \pm 93 dolphins). A summary of the sampling effort and data gathered throughout the surveys is presented in Table 1.

Overall, 462 Guiana dolphins were identified and catalogued based on markings found on their dorsal fins. There was no correlation between the number of animals captured in each survey and the cumulative number of captured individuals ($r = -0.12$, $P = 0.66$; $N = 17$). However, there was a significant positive relationship between the number of animals recaptured in each survey and the cumulative number of captured individuals ($r = 0.66$; $P = 0.003$; $N = 17$). This pattern was also observed when calculating recapture rates (number of recaptured individuals divided by the cumulative number of captured animals) and capture rates (number of captured individuals divided by the cumulative number of




Fig. 1. Map of the study area showing the geographical position of *Sotalia guianensis* groups (+) at the delimited area of western Ilha Grande Bay, south-eastern Brazil.

captured animals) (Figure 2). The number of animals recaptured in each survey ranged from 0 to 79 (mean 39.4 ± 23.6 animals).

In Ilha Grande Bay, 167 Guiana dolphins (36.1%) were sighted only once and were regarded as non-resident dolphins (Table 2). The other 295 dolphins (63.9%) were seen two or more times and were therefore considered for evaluation of their degrees of residence in the study area. The number of times these dolphins were sighted ranged from 2 to 11 (Table 2). The time interval between the first and last sightings of any one dolphin ranged from 2 to 310 days, while the majority of the dolphins ($N = 111$; 37.6%) exhibited maximum residence periods of between 93 and 118 days

(Figure 3). The average number of days between the first and last sightings was 121 days (± 99.1 days).

The average number of days between adjacent sightings for an individual was 35 days (± 27.4 days), with durations ranging from 1 to 153 days. Some of the resident dolphins were sighted between 47 and 105 days ($N = 113$; 38.3%), while 99 dolphins (33.6%) had mean intervals of 17 and 44 days. Only one dolphin (0.3%) had a mean interval between resightings up to 150 days. Eighty-two animals (27.8%) were sighted at mean intervals ranging from 1 to 4 days.

After considering all three parameters together, we were able to determine how many *Sotalia guianensis* dolphins had high, medium and low degrees of residence in Ilha

Table 1. Summary of sample effort and data gathered during the study period in Ilha Grande Bay.

Boat survey date	Group size	Mean subgroup size	Number of photographs taken	Number of new identifications	Number of resightings
5 May 2007	60	10.1	831	26	0
7 May 2007	200	10.2	1488	61	16
13 May 2007	80	9.2	283	6	13
25 August 2007	200	9.4	807	35	30
26 August 2007	150	11.9	1219	22	55
27 August 2007	100	12.9	888	24	53
29 August 2007	20	10.9	186	1	11
30 August 2007	300	22.2	1280	60	20
30 November 2007	150	24.9	465	8	30
1 December 2007	350	15.4	2821	96	56
2 December 2007	300	19.3	1788	27	79
3 December 2007	80	10.8	691	15	52
5 March 2008	200	32	1133	29	52
6 March 2008	150	13.3	1017	8	27
7 March 2008	100	13.4	1251	23	69
8 March 2008	250	27.1	1137	17	73
9 March 2008	120	16.1	684	4	34




Fig. 2. Capture and recapture rates of Guiana dolphins in Ilha Grande Bay. From 2 December to 9 March the proportions of recaptured animals were higher than the capture rates.

Grande Bay (Figure 4). The reference numbers used for each parameter and degree of residence are presented in Table 3. In summary, 36.1% of the dolphins (N = 167) had null residence as they were not recaptured; 17.7% (N = 82 dolphins) had a low degree of residence; 24.8% (N = 114 dolphins) had a medium degree of residence; and 21.4% (N = 99 dolphins) had a high degree of residence in Ilha Grande Bay.

DISCUSSION

The number of identified animals levelled off with time, suggesting that most of the dolphins in Ilha Grande Bay were identified. Recapture rates tended to be higher than capture rates on the latest surveys and the correlation between the number of recaptures and the cumulative curve of identified individuals was positive. Additionally, the number of recaptures tended to increase during the study period and the number of new identifications tended to decrease with time. For this reason, we believe that our sample effort was representative for this population.

As stated by Ballance (1990), each measure of residence must be interpreted with caution. Some dolphins were frequently sighted on consecutive days. An animal that visited Ilha Grande Bay only a few times on consecutive days would have a high number of resightings yet a relatively low

Table 2. Frequencies of non-resident and resident Guiana dolphins. A total of 295 dolphins were used to establish degrees of residency within the population. None of the dolphins were sighted 8 or 10 times.

	Number of sightings	Number of individuals	Frequency (%)
Non-residents	1	167	36.1
Residents	2	118	25.5
	3	73	15.8
	4	49	10.6
	5	29	6.3
	6	19	4.2
	7	5	1.1
	9	1	0.2
	11	1	0.2




Fig. 3. Maximum residence periods of resident Guiana dolphins in Ilha Grande Bay. The average number of days between first and last sightings was 121 days (± 99.1 days).

degree of residence. Alternatively, an animal may have been frequently sighted over an extended period of up to 90 days. This would indicate a high degree of residence. The average number of days between sightings takes into account both the number of times an animal was sighted and the time between its first and last sightings. Individuals that visited Ilha Grande Bay on only a few occasions spaced widely apart in time exhibited a medium degree of residence. In general, Guiana dolphins exhibited varying degrees of residence in Ilha Grande Bay. The results provided by this study indicate that some individuals (46%) frequently used the bay during the sample period, exhibiting high and medium degrees of residence. Meanwhile, the majority of photo-identified dolphins (54%), namely low resident and non-resident dolphins, were less frequent. This fact suggests that this population has a fluid structure as different individuals were observed to compose it during the study period. The large proportion of dolphins seen only once indicates that many individuals spend most of their time outside of the study area, either offshore or in adjacent waters (Parra *et al.*, 2006). Similar environmental features have been found for




Fig. 4. Scatterplot of maximum residence periods versus the mean number of days, showing distinct degrees of residence for Guiana dolphins in Ilha Grande Bay. High level = 99 dolphins; medium level = 114 dolphins; low level = 82 dolphins.

Table 3. Reference numbers used for each parameter according to each degree of residence.

Degree of residence	Number of resightings	Number of days between first and last sightings	Mean number of days between resightings
Low	2–4	2–7	1–3.5
Medium	2–6	93–310	46.5–153
High	3–11	94–308	16.5–44

the *S. guianensis* population of Sepetiba Bay, which is 70 km from the study area. Nery *et al.* (2008) found that 64% of the identified dolphins were seen only one time, suggesting that dolphins inhabiting Sepetiba Bay may also use Ilha Grande Bay as it offers resources and shelter against natural predators. Thus, Ilha Grande Bay appears to be an important part of the home range of Guiana dolphins on the southern coast of Rio de Janeiro State. Further comparisons between both photo-identification catalogues would be necessary to answer this question.

The absence of a standard method for residence analysis makes a comparison of this study's results to others' difficult. The main difference is related to the time scale considered. Several studies have observed that Guiana dolphins may show long-term residency in particular areas along the Brazilian coast (Azevedo *et al.*, 2004; Flores & Bazzalo, 2004; Rossi-Santos *et al.*, 2007; Nery *et al.*, 2008). In Sepetiba Bay, for instance, one dolphin remained in the area for over 11 years (Nery *et al.*, 2008). Our study draws conclusions regarding residency based on days instead of years. However, Ballance (1990) gathered information about residency of bottlenose dolphins (*Tursiops truncatus*) in Kino Bay, Mexico, based on observations made during an eight-month study. A continuum of survey efforts in Ilha Grande Bay could be necessary to determine if a similar pattern to that already described for *S. guianensis* occurs.

Abundance, distribution and availability of resources in a certain habitat determine the home range size of a species (Ballance, 1992). Marine environments may be composed of a variety of microhabitats, each of them providing different resources for individuals in a population. Ilha Grande Bay presents a great diversity of microhabitats, which enhances local productivity (Nogara, 2000). Data about annual and spatial distributions and abundances of food resources (mainly fish, cephalopods and shrimps) are scarce for Ilha Grande Bay. However, many prey items described for Guiana dolphins are found in Ilha Grande Bay, such as squid (*Loligo plei*, *L. sanpaulensis* and *Lolliguncula brevis*) and demersal and pelagic fish (*Trichiurus lepturus*, *Centropomus* sp., *Pomatomus saltatrix*, *Cynoscion leiarchus*, *C. striatus* and *C. jamaicensis*) (Lodi, 2003a). A previous study of habitat use by Guiana dolphins in Ilha Grande Bay proposed that the area is used mainly for foraging and feeding purposes (Lodi, 2003a).

The presence of the dolphins in the area may be a response to their abundance and distribution of prey. In Sepetiba Bay, it was reported that fish communities vary according to the season of the year (Araújo *et al.*, 1998). Some fish species showed habitat preferences, others had a patchy distribution around the bay, and still others could be found year-round (Araújo *et al.*, 1998). We believe that such fluctuations in prey communities have a strong effect on the Guiana dolphins

of Ilha Grande Bay. When food resources display a patchy distribution in the bay, dolphins may present a similar pattern to optimize their individual food intake. Delphinids spend considerable time and energy in locating, pursuing, capturing and processing prey (Williams *et al.*, 2001). As the energy costs of living in water are high, they must acquire enough energy from the intake of prey. Wedekin *et al.* (2007) observed that Guiana dolphins in Norte Bay, southern Brazil, use a greater area during months with low resource availability (from March to September). It seems plausible that Guiana dolphins' ranges in Ilha Grande Bay extended beyond the area covered in this study, which may have affected individuals' residency identification.

Estuarine waters are an important breeding and calving area for Guiana dolphins (Santos *et al.*, 2001). The high proportion of calves and juveniles reported by the present study (45%) may be related to local resource availability. A similar proportion was described for Sepetiba Bay, south-eastern Brazil (47%) (Nery, 2008). In Guanabara Bay, also in south-eastern Brazil, young animals corresponded to 26% of the total photographs taken (Azevedo, 2005). Lodi (2003b) found a high proportion of groups with calves in Ilha Grande Bay (88.9%), suggesting that it may represent an important calving area because of its sheltered waters (without strong winds and currents), absence of natural predators and prey availability. Besides, large groups are reported to live in the area. This may provide protection to newborns and increase opportunities for learning and development (Wells *et al.*, 1987). Thus, females with newborns would have a place for teaching and raising their calves without major threats to their safety.

Ilha Grande Bay is regarded as a priority area for marine conservation by the Brazilian Environmental Agency (IBAMA) because of the high levels of biodiversity described in the area (Creed *et al.*, 2007). Our findings suggest that this bay represents an important habitat for *Sotalia guianensis* dolphins as well. Preliminary abundance estimates have already been provided for this population, taking into consideration an overlap among confidence intervals of three different estimators for closed-population models (Espécie *et al.*, 2009). The results have indicated that Ilha Grande Bay holds the biggest population of *Sotalia guianensis* in its entire geographical distribution, ranging from 1041 to 1142 individuals (Espécie *et al.*, 2009). Conclusions based on short-term data tend to be transitory and simplistic (Scott *et al.*, 1990). However, the results presented here provide the first report of the residence dynamics of Guiana dolphins in Ilha Grande Bay, showing that the area is regularly used by both adults and young individuals. The recurrent use of Ilha Grande Bay by some dolphins raises some concern about the long-term survival of local populations if the current habitat quality is not improved or at least maintained. Species with high levels of site fidelity and residency are vulnerable to population declines due to habitat degradation and loss (Warkentin & Hernandez, 1996). According to Nery *et al.* (2008), anthropogenic influences have been rapidly increasing in Sepetiba Bay due to its economic interest for Rio de Janeiro State, and Ilha Grande Bay may become a refuge for *Sotalia guianensis* when conditions worsen in Sepetiba Bay. Like many other coastal areas, Ilha Grande Bay is also suffering from intense economic development with little or no consideration for marine habitats and species. For this reason, further investigations into many

ecological parameters of this *S. guianensis* population are still necessary for designing a conservation plan for this species in the area.

ACKNOWLEDGEMENTS

This project was made possible by the logistical support of Instituto Aqualie. We extend our thanks to Bill Rossiter (Cetacean Society International). We appreciate the comments and suggestions from Carlos Eduardo L. Esbérard and two anonymous referees of manuscript drafts. Our sincere thanks go to Sergio C. Moreira for preparing the map. Our deepest gratitude goes to Lauren Stanton for reviewing the English version of this paper. We would also like to thank Dona Elsa for housing us during data collection, Gilberto and Tico for their friendship and invaluable skills with the survey boat, and many trainee students from Laboratório de Bioacústica e Ecologia de Cetáceos, Universidade Federal Rural do Rio de Janeiro. This research was supported by Cetacean Society International (M.A.E., grant number 659-622-6826); the Conselho Nacional de Desenvolvimento Científico e Tecnológico (M.A.E., grant number 111555/2008-6); and the Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (R.H.O.T., grant number E-26/151.047/2007).

REFERENCES

- Araújo F.G., Cruz-Filho A.G., Azevêdo M.C.C. and Santos A.C.A. (1998) Estrutura da comunidade de peixes demersais da Baía de Sepetiba, RJ. *Revista Brasileira de Biologia* 58, 417–430.
- Azevedo A.F. (2005) *Comportamento e uso do espaço por Sotalia guianensis (Cetacea, Delphinidae) na Baía de Guanabara (RJ) e variação geográfica dos assobios da espécie ao longo da costa do Brasil*. DSc thesis. Universidade Estadual do Rio de Janeiro, Rio de Janeiro, Brazil.
- Azevedo A.F., Lailson-Brito J. Jr, Cunha H.A. and Van Sluys M. (2004) Site fidelity of marine tucuxis (*Sotalia fluviatilis*) in Guanabara Bay, south-eastern Brazil. *Journal of Cetacean Research and Management* 6, 265–268.
- Ballance L.T. (1990) Residence patterns, group organization and surface association of bottlenose dolphins in Kino Bay, Gulf of California, Mexico. In Leatherwood S. and Reeves R.R. (eds) *The bottlenose dolphin*. San Diego: Academic Press, pp. 267–284.
- Ballance L.T. (1992) Habitat use patterns and ranges of the bottlenose dolphin in the Gulf of California, Mexico. *Marine Mammal Science* 8, 262–274.
- Creed J.C., Absalão R.S., Brasil A.C.S., Figueiredo M.A.O., Ferreira C.E.L., Pires D.O., Serejo C.S. and Ventura C.R.R. (2007) Sumário executivo. In Creed J.C., Pires D.O. and Figueiredo M.A.O. (eds) *Biodiversidade marinha da Baía da Ilha Grande*. Brasília: Ministério do Meio Ambiente, pp. 19–39.
- Espécie M.A., Kiatkowski D.A., Tardin R.H.O. and Simão S.M. (2009) Abundance estimates of Guiana dolphins (*Sotalia guianensis*) in Ilha Grande Bay, Brazil. In Nascimento E. (ed.) *Anais do III Congresso Latino Americano de Ecologia, Hotel Guanabara, São Lourenço, 10–13 September 2009. Ecologia marinha*. São Paulo: Imprensa da Universidade de São Paulo, pp. 1–3.
- Flores P.A.C. (1999) Preliminary results of a photo-identification study of the marine tucuxi *Sotalia fluviatilis* in southern Brazil. *Marine Mammal Science* 15, 840–847.
- Flores P.A.C. and Bazzalo M. (2004) Home range and movement patterns of the marine tucuxi, *Sotalia fluviatilis*, in Baía Norte, southern Brazil. *Latin American Journal of Aquatic Mammals* 3, 37–52.
- Gunnlaugsson T. and Sigurjónsson J. (1990) A note on the problem of false positives in the use of natural marking data and abundance estimation. In Hammond P.S., Mizroch S.A. and Donovan G.P. (eds) *Individual recognition of cetaceans: use of photoidentification and other techniques to estimate population parameters*. Cambridge: International Whaling Commission, Special Issue 12, pp. 143–145.
- Hammond P.S., Mizroch S.A. and Donovan G.P. (1990) *Individual recognition of cetaceans: use of photoidentification and other techniques to estimate population parameters*. Special Issue 12. Cambridge: International Whaling Commission.
- Lodi L. (2003a) Seleção e uso de habitat pelo boto-cinza, *Sotalia guianensis* (van Béneden, 1864) (Cetacea, Delphinidae), na Baía de Paraty, estado do Rio de Janeiro. *Bioikos* 17, 5–20.
- Lodi L. (2003b) Tamanho e composição de grupo dos botos-cinza, *Sotalia guianensis* (Van Béneden, 1864) (Cetacea, Delphinidae), na Baía de Paraty, Rio de Janeiro, Brasil. *Atlântica* 25, 135–146.
- Lodi L. and Hetzel B. (1998) Grandes agregações do boto-cinza (*Sotalia fluviatilis*) na Baía da Ilha Grande, Rio de Janeiro. *Bioikos* 12, 26–30.
- Mahiques M.M. (1987) *Considerações sobre os sedimentos de superfície de fundo da Baía da Ilha Grande, estado do Rio de Janeiro*. MSc dissertation. Universidade de São Paulo, São Paulo, Brazil.
- Mazzoil M., McCulloch S.D., Defran R.H. and Murdoch E.M. (2004) Use of digital photography and analysis of dorsal fins for photoidentification of bottlenose dolphin. *Aquatic Mammals* 30, 209–210.
- Mazzoil M., Reif J.S., Youngbluth M., Murdoch E., Bechdel S.E., Howells E., McCulloch S.D., Hansen L.J. and Bossart G.D. (2008) Home ranges of bottlenose dolphins (*Tursiops truncatus*) in the Indian River Lagoon, Florida: environmental correlates and implications for management strategies. *EcoHealth* 5, 278–288.
- Nery M.F. (2008) *Fidelidade de habitat e estimativa populacional de Sotalia guianensis (Cetacea, Delphinidae) da Baía de Sepetiba—RJ, por meio da técnica de marcação-recaptura*. MSc dissertation. Universidade Federal Rural do Rio de Janeiro, Seropédica, Brazil.
- Nery M.F., Espécie M.A. and Simão S.M. (2008) Site fidelity of *Sotalia guianensis* (Cetacea, Delphinidae) in Sepetiba Bay, Rio de Janeiro, Brazil. *Revista Brasileira de Zoologia* 25, 182–187.
- Nogara P.J. (2000) *Caracterização dos ambientes marinhos da Área de Proteção Ambiental de Cairuçu—Município de Paraty—RJ*. Fundação SOS Mata Atlântica (Technical report), 83 pp.
- Parra G.J., Corkeron P.J. and Marsh H. (2006) Population sizes, site fidelity and residence patterns of Australian snubfin and Indo-Pacific humpback dolphins: implications for conservation. *Biological Conservation* 129, 167–180.
- Pizzorno J.L.A. (1999) *Estimativa populacional do boto-cinza, Sotalia fluviatilis, na Baía de Guanabara, por meio de fotoidentificação*. MSc dissertation. Universidade Federal Rural do Rio de Janeiro, Seropédica, Brazil.
- Rossi-Santos M.R., Wedekin L.L. and Monteiro-Filho E.L.A. (2007) Residence and site fidelity of *Sotalia guianensis* in the Caravelas River estuary, eastern Brazil. *Journal of the Marine Biological Association of the United Kingdom* 87, 207–212.
- Santos M.C.O., Acuña L.B. and Rosso S. (2001) Insights on site fidelity and calving intervals of the marine tucuxi dolphin (*Sotalia fluviatilis*) in south-eastern Brazil. *Journal of the Marine Biological Association of the United Kingdom* 81, 1049–1052.
- Scott M.D., Wells R.S. and Irvine A.B. (1990) A long term study of bottlenose dolphins on the west coast of Florida. In Leatherwood S. and

- Reeves R.R. (eds) *The bottlenose dolphin*. San Diego: Academic Press, pp. 235–244.
- Signorini S.R.** (1980) A study of the circulation in bay of Ilha Grande and bay of Sepetiba part I, an assessment of the tidally and wind-driven circulation using a finite element numerical model. *Boletim do Instituto Oceanográfico* 29, 41–55.
- Simão S.M., Pizzorno J.L.A., Perry V.N. and Siciliano S.** (2000) Aplicação da técnica de fotoidentificação do boto-cinza, *Sotalia fluviatilis* (Cetacea, Delphinidae), da Baía de Sepetiba. *Floresta e Ambiente* 7, 31–39.
- Stevick P.T., Palsboll P.J., Smith T.D., Bravington M.V. and Hammond P.S.** (2001) Errors in identification using natural markings: rates, sources, and effects on capture–recapture estimates of abundance. *Canadian Journal of Fisheries and Aquaculture Science* 58, 1861–1870.
- Warkentin I.G. and Hernandez D.** (1996) The conservation implications of site fidelity: a case-study involving nearctic–neotropical migrant songbirds wintering a Costa Rican mangrove. *Biological Conservation* 77, 143–150.
- Wedekin L.L., Daura-Jorge F.G., Piacentini V.Q. and Simões-Lopes P.C.** (2007) Seasonal variations in spatial usage by the estuarine dolphin, *Sotalia guianensis* (van Bénédén, 1864) (Cetacea; Delphinidae) at its southern limit of distribution. *Brazilian Journal of Biology* 67, 1–8.
- Wells R.S.** (1991) The role of long-term study in understanding the social structure of a bottlenose dolphin community. In Pryor K. and Norris K.S. (eds) *Dolphins societies: discoveries and puzzles*. Berkeley: University of California Press, pp. 199–225.
- Wells R.S. and Scott M.D.** (1990) Estimating bottlenose dolphin population parameters from individual identification and capture–release techniques. In Hammond P.S., Mizroch S.A. and Donovan G.P. (eds) *Individual recognition of cetaceans: use of photoidentification and other techniques to estimate population parameters*. Cambridge: International Whaling Commission, Special Issue 12, pp. 470–416.
- Wells R.S., Scott M.D. and Irvine A.B.** (1987) The social structure of free-ranging bottlenose dolphins. In Genoways H.H. (ed.) *Current mammalogy*. New York: Plenum Press, pp. 247–305.
- Williams T.M., Haun J., Davis R.W., Fuiman L.A. and Kohin S.** (2001) A killer appetite: metabolic consequences of carnivory in marine mammals. *Comparative Biochemistry and Physiology Part A* 129, 785–796.
- and
- Würsig B. and Würsig M.** (1977) The photographic determination of group size, composition and stability of coastal porpoises (*Tursiops truncatus*). *Science* 198, 755–756.

Correspondence should be addressed to:

M.A. Espécie
 Laboratório de Bioacústica e Ecologia de Cetáceos
 Departamento de Ciências Ambientais
 Universidade Federal Rural do Rio de Janeiro
 Rodovia BR 465, km 7 Seropédica
 Rio de Janeiro, 23890-000, Brazil
 email: mariana_especie@ufrj.br